

---

EXPERIMENTAL  
ARTICLES

---

## Screening of Marine Bacteria for Fucoidanases

I. Yu. Bakunina\*, L. S. Shevchenko\*, O. I. Nedashkovskaya\*,  
N. M. Shevchenko\*, S. A. Alekseeva\*\*, V. V. Mikhailov\*, and T. N. Zvyagintseva\*

\*Pacific Institute of Bioorganic Chemistry, Far Eastern Division,  
Russian Academy of Sciences, pr. 100-letiya Vladivostoka 159, Vladivostok, 690022 Russia

\*\*Far Eastern State University, ul. Sukhanova 8, Vladivostok, 690600 Russia

Received June 29, 1999

**Abstract**—Twenty-five strains of epiphytic marine bacteria isolated from the brown algae *Fucus evanescens* and *Chorda filum* and fifty-three bacteria isolated from the sea cucumber *Apostichopus japonicus* were screened for fucoidanases using fucoidans prepared from the brown algae *F. evanescens*, *Laminaria cichorioides*, and *L. japonica*. Eighteen bacterial epiphytes and thirty-eight bacterial isolates from the sea cucumber were found to contain fucoidanases, which were able to hydrolyze either all of the fucoidans studied or some of them. Bacteria of the genera *Cytophaga* and *Aalteromonas/Pseudoalteromonas* exhibited the highest fucoidanase activities, which, however, did not exceed the activity of fucoidanases from the already known sources.

**Key words:** marine bacteria, brown algae, sea cucumbers, fucoidans, fucoidanases

Fucoidans are widespread among marine macrophytes. In brown algae, they primarily play the role of the cell-wall matrix polysaccharides [1]. Fucoidans have also been found in the envelope of sea cucumbers [2]. In sea urchins, they are involved in the sperm-egg adhesion [3]. Fucoidans exhibit a wide range of activities toward different mammalian organs [3–5]. The biological activity of fucoidans is evidently determined by their specific chemical structure: fucoidans represent a family of homo- and heteropolysaccharides composed mainly of fucose residues sulfated at positions 2 and/or 4 and bound by  $\alpha$ -1,2- or  $\alpha$ -1,3-*O*-glycosidic bonds [6, 7]. In addition to fucose, fucoidans may also contain mannose, xylose, galactose, uronic acids, and rhamnose. The detailed chemical structure of fucoidans remains unknown.

Investigation of fucoidanases may provide insight into the structure and mechanism of the biological activity of fucoidans. Data available in the literature mainly concern the fucoidanases of molluscs [8, 9] and echinoderms [10], whereas information on microbial fucoidanases is scarce [11].

The aim of the present work was to screen marine bacteria for the efficient producers of fucoidanases.

### MATERIALS AND METHODS

**Microorganisms and cultivation conditions.** The isolates of marine bacteria studied in the present work were obtained during scientific offshore expeditions aboard the research vessel *Academician Oparin* in the Sea of Okhotsk near the Kuril islands of Paramushir, Onekotan, and Iturup, as well as at the Marine Experi-

mental Station of the Pacific Institute of Bioorganic Chemistry (PIBOC) in the Peter the Great Bay. At present, these isolates are stored in the PIBOC Collection of Marine Microorganisms (PIBOC CMM).

The isolation and identification of marine bacteria from the brown algae *Fucus evanescens* and *Chorda filum* and the sea cucumber *Apostichopus japonicus* were described elsewhere [12].

Bacterial strains for the primary screening of fucoidanase producers were grown on a shaker (160 rpm) at 28°C for 24 h in a medium containing (g/l sea water) Difco bacto-peptone, 5.0; Difco yeast extract, 2.0; fucoidan 1.0; K<sub>2</sub>HPO<sub>4</sub>, 0.2; and MgSO<sub>4</sub>, 0.05 (pH 7.5–7.8).

**Preparation of bacterial extracts.** Bacterial cells (0.5 g) were suspended in 1 ml of 0.05 M phosphate buffer (pH 7.2) and disrupted by sonication in a UZDN-2 ultrasonic disintegrator. The cell homogenate was centrifuged at 10000 g for 30 min, and the supernatant was assayed for fucoidanase activity using fucoidans F<sub>1</sub>, F<sub>2</sub>, and F<sub>3</sub> prepared from the brown algae *F. evanescens*, *Laminaria cichorioides*, and *L. japonica*, respectively [13].

**Isolation and characterization of fucoidans.** Ground wet or frozen seaweeds were treated with ethanol and acetone and then extracted with 0.4% HCl at 20–25°C (cold extraction) and with hot water at 60–70°C (hot extraction). The cold- and hot- extracted polysaccharides were subjected to hydrophobic chromatography. The polysaccharides adsorbed on hydrophobic adsorbent were eluted with water and then with a gradient of ethanol in water. Fucoidans that were

**Table 1.** Some relevant characteristics of fucoidans used as enzyme substrates

Fucoidan	Carbohydrate content, % of dry wt	Molecular mass, kDa	Carbohydrate composition, %	Fuc/SO <sub>4</sub> mol. ratio	IR(SO <sub>4</sub> ) maximum, cm <sup>-1</sup>
F <sub>1</sub> from <i>F. evanescens</i>	28	300–500	Fuc, Gal, Xyl, 87 : 1 : 10	1 : 0.8	820
F <sub>2</sub> from <i>L. cichorioides</i>	30	60–80	Fuc, Gal, Xyl, GlcUa 82 : 10 : 2 : 5	1 : 1.7	842
F <sub>3</sub> from <i>L. japonica</i>	22	28–30	Fuc, Gal, Man, Xyl 42 : 40 : 9 : 2	1 : 0.9	842

**Table 2.** Specific fucoidanase activity\* of marine bacteria isolated from the brown alga *F. evanescens*

Taxonomic position of strain	Strain	Fucoidan used as enzyme substrate		
		F <sub>1</sub> from <i>F. evanescens</i>	F <sub>2</sub> from <i>L. cichorioides</i>	F <sub>3</sub> from <i>L. japonica</i>
<i>Cytophaga</i> sp.	1F2	0	0	3510
<i>Cytophaga</i> sp.	2F3	0	0	2980
<i>Cytophaga</i> sp.	2F5	0	0	2190
<i>Cytophaga</i> sp.	2F7	0	4000	4000
<i>Cytophaga</i> sp.	2F9B	660	4980	1010
<i>Cytophaga</i> sp.	2F13	1140	12550	2140
<i>Cytophaga</i> sp.	2F16	600	4120	1520
<i>Cytophaga</i> sp.	12F2	0	6220	0
<i>Cytophaga</i> sp.	12F5	0	0	17680
<i>Cytophaga</i> sp.	12F9	0	8590	8590
<i>Flexibacter/Cytophaga</i>	12F8	3920	9650	4130
<i>Flexibacter</i> sp.	12F6	0	0	6710
<i>Pseudoalteromonas</i> sp.	2F10	710	12090	37360
<i>Pseudoalteromonas</i> sp.	12F1	220	8600	0
<i>Pseudoalteromonas</i> sp.	12F3	370	8680	0
<i>Pseudoalteromonas</i> sp.	1F5	1490	19650	8940
<i>Pseudoalteromonas</i> sp.	20-92	5290	3640	–
<i>Pseudoalteromonas</i> sp.	20-101-1**	2130	2630	–

\* Specific activity was expressed as enzymatic units per 1 g of wet bacterial biomass.

\*\* This strain was isolated from the brown alga *Chorda filum*.

eluted with water were precipitated with 80% ethanol and reprecipitated with 60% ethanol.

The molecular mass of fucoidans was determined by gel filtration on Sephadex G-50 and Sepharose CL-4B columns (100 × 1 cm; water, 15 ml/h) using 10-, 20-, 40-, and 80-kDa dextrans as molecular weight markers. Detection of polysaccharides in fractions was performed colorimetrically with the phenol–sulfuric acid reagent [14].

The carbohydrate composition of fucoidans was determined after hydrolyzing them in 4 N HCl at 100°C for 2 h. The hydrolysates were analyzed on a Biotronik carbohydrate analyzer (0.63 × 30 cm Durrum X4-20 column; 60°C; bicinchoninic acid reagent; Shimadzu CR 2AX detector) using the authentic samples of fucose (Fuc), glucose (Glu), galactose (Gal), mannose (Man), xylose (Xyl), and rhamnose (Rha) purchased from Serva (Germany) as reference standards.

Sulfate groups were determined turbidimetrically after the hydrolysis of fucoidans in 4 N HCl and the addition of a BaCl<sub>2</sub> suspension in gelatin. The IR spectra of fucoidans were recorded on a Carl Zeiss IR-75 spectrophotometer. Some relevant characteristics of the fucoidans used in this study are presented in Table 1.

**Assay of the fucoidanase activity of bacterial cell extracts.** A solution (0.4 ml) containing 4 mg of fucoidan in 0.05 M phosphate buffer (pH 7.2) was mixed with 0.1 ml of bacterial cell extract and incubated at 20°C for 20 h. Fucoidanase activity was estimated from the increase in the reducing sugar content of the reaction mixture. The concentration of reducing sugars was determined by the Nelson–Somogyi method [15]. One unit of fucoidanase activity was defined as the amount of enzyme releasing 1 nmole of fucose from fucoidan in 20 h.

**Table 3.** Fucoidanase activity of marine bacteria isolated from the sea cucumber *A. japonicus*

Taxonomic position of strain	Strain	Fucoidan used as enzyme substrate	
		F <sub>1</sub> from <i>F. evanescens</i>	F <sub>2</sub> from <i>L. cichorioides</i>
<i>Acinetobacter</i> sp.	M-3HL-8/1	0	240
<i>Alteromonas/Pseudoalteromonas</i>	M-3HB-8/1	1150	230
<i>Alteromonas/Pseudoalteromonas</i>	M-3HB-8/2	40	1270
<i>Alteromonas/Pseudoalteromonas</i>	M-3HB-9	420	2940
<i>Alteromonas/Pseudoalteromonas</i>	M-3HB-15/3	1540	5300
<i>Alteromonas/Pseudoalteromonas</i>	M-3HB-17/1	330	1260
<i>Alteromonas/Pseudoalteromonas</i>	M-3HB-18	1010	3150
<i>Alteromonas/Pseudoalteromonas</i>	M-3HB-19	0	1530
<i>Alteromonas/Pseudoalteromonas</i>	M-3HL-6/2	3180	1060
<i>Alteromonas/Pseudoalteromonas</i>	M-3HL-14/2	2420	0
<i>Alteromonas/Pseudoalteromonas</i>	M-3HL-14/3	1520	450
<i>Alteromonas/Pseudoalteromonas</i>	M-3HL-18/1	0	2000
<i>Alteromonas/Pseudoalteromonas</i>	M-3HL-21/1	220	60
<i>Bacillus</i> sp.	M-3HL-1/2	5790	2320
Coryneforms	M-3HB-14/2	370	2280
Coryneforms	M-3HB-25/2	930	2190
Coryneforms	M-3HB-27/1	2010	1590
Coryneforms	M-3HB-27/2	0	810
Coryneforms	M-3HL-5/2	1520	0
<i>Cytophaga</i> sp.	M-3HB-28/1	170	120
<i>Flavobacterium</i> sp.	M-3HB-26	0	1040
<i>Halomonas</i> sp.	M-3HL-13/2	990	610
<i>Halomonas</i> sp.	M-3HL-17/3	330	910
<i>Pseudoalteromonas</i> sp.	M-3HB-2	2230	1480
<i>Pseudomonas/Halomonas</i>	M-3HB-14/1	1230	4960
<i>Pseudomonas/Halomonas</i>	M-3HB-15/2	0	1310
<i>Pseudomonas</i> sp.	M-3HL-7/1	3060	2350
<i>Pseudomonas</i> sp.	M-3HL-9	1470	680
<i>Micrococcus</i> sp.	M-3HB-15/1	720	1980
<i>Micrococcus</i> sp.	M-3HB-25/3	0	1540
<i>Vibrio</i> sp.	M-3HB-3	450	1230
<i>Vibrio</i> sp.	M-3HB-11/2	970	1700
<i>Vibrio</i> sp.	M-3HL-6/3	0	120
Unknown	M-3HL-2	3360	3950
Unknown	M-3HL-7/2	3350	340
Unknown	M-3HB-15/4	660	950
Unknown	M-3HB-16/1	0	340
Unknown	M-3HB-16/2	440	630

## RESULTS AND DISCUSSION

Marine microorganisms are promising sources of valuable biologically active compounds, including rare enzymes [16, 17]. Some marine bacteria are able to

degrade the insoluble polysaccharides chitin and agar, as well as the cell-wall polysaccharides of seaweeds [18] and other organisms. When damaged during storms, brown seaweeds become the subject of attack by epiphytic bacteria, which possess the necessary

**Table 4.** Taxonomic composition of fucooidan-degrading marine bacteria

Taxon	Number of strains isolated from					
	sea cucumber		sea alga		total	
	all	active	all	active	all	active
<i>Acinetobacter</i>	1	1	–	–	1	1
<i>Alteromonas/Pseudoalteromonas</i>	23	13	8	6	31	19
<i>Bacillus</i>	1	1	–	–	1	1
Coryneforms	6	4	–	–	6	4
<i>Cytophaga/Flexibacter</i>	2	1	12	12	14	13
<i>Flavobacterium</i>	1	1	–	–	1	1
<i>Micrococcus</i>	4	3	1	0	5	3
<i>Pseudomonas/Halomonas</i>	6	6	–	–	6	6
<i>Vibrio</i>	4	3	2	0	6	3
Unidentified	6	5	2	0	8	2
Total	53	38	25	18	78	56

enzymatic systems capable of degrading algal cell-wall polysaccharides to mono- and oligosaccharides. These saccharides are utilized not only by epiphytic bacteria themselves, but also by other marine microorganisms. Glycosidases are often produced by symbiotrophic bacteria. For instance, bacteria of the genus *Vibrio* associated with the brown alga *Laminaria longicruris* are able to hydrolyze laminarin [19], whereas bacteria isolated from the red algae *Polisiphonia lanosa*, *Hypnea charoides*, and *Gracilaria gracilis* are able to hydrolyze agar [20, 21].

These data suggest that the seaweeds under study can also be a source of nutrients for bacterial epiphytes. On the other hand, the envelope of sea cucumbers contains fucooidans [2]; therefore, the associated bacteria may possess enzymes capable of degrading these polysaccharides. For these reasons, screening for fucooidanases was primarily carried out among the marine bacteria isolated from the brown alga *F. evanescens* (Table 2) and from the coelomic liquid and homogenates of the sea cucumber *A. japonicus* (Table 3), which is abundant in the Peter the Great Bay of the Sea of Japan.

Twenty-three of the twenty-five strains of heterotrophic bacteria isolated from the brown algae *F. evanescens* and *Chorda filum* were identified to the genus level (Table 3). Similar to other marine bacteria [22], the majority of the epiphytes isolated turned out to be gram-negative aerobic rods.

All of the strains studied, except 20-92 and 20-101/1, were analyzed for the presence of fucooidanases using the hot-extracted fucooidans F<sub>1</sub>, F<sub>2</sub>, and F<sub>3</sub> of the brown algae *F. evanescens*, *L. cichorioides*, and *L. japonica*. Eighteen of the twenty-five bacteria associated with the brown seaweeds were able to hydrolyze fucooidans

(Table 4). All of the bacteria of the genus *Cytophaga* under study exhibited fucooidanase activity (Table 2). Five of the twelve bacterial strains of this genus produced fucooidanases that were able to hydrolyze only fucooidan F<sub>3</sub>. The fucooidanase of *Cytophaga* sp. 12F2 was specific for fucooidan F<sub>2</sub>, and the fucooidanases of *Cytophaga* sp. 2F7 and 12F9 were specific for fucooidans F<sub>2</sub> and F<sub>3</sub>. The fucooidanases of *Cytophaga* spp. 2F9B, 2F13, 2F16, and 12F8 exhibited the widest substrate specificity: they were able to degrade all three fucooidans under study.

Six of the eight bacterial epiphytes of the genus *Pseudoalteromonas* also possessed fucooidanase activity (Table 2). The fucooidanase of *Pseudoalteromonas* sp. 12F1 was able to hydrolyze only fucooidan F<sub>3</sub>, whereas the fucooidanases of other strains of this species could hydrolyze all three fucooidans (it should be noted that the fucooidanase activity of these strains was higher than that of the fucooidan-degrading *Cytophaga* sp. None of the bacterial strains belonging to the genera *Vibrio* and *Micrococcus* could degrade fucooidans. Fifty-three bacterial strains associated with the sea cucumber *A. japonicus* were tested for fucooidanase activity using two fucooidans, F<sub>1</sub> and F<sub>2</sub> (Table 3). This bacterial group was dominated by gram-negative aerobic rodlike bacteria of the genera *Alteromonas/Pseudoalteromonas*. Some fucooidanase-producing bacteria of this group belonged to coryneforms and to the genera *Vibrio*, *Micrococcus*, and *Halomonas* (Tables 3 and 4). The thirty-eight bacterial associates of *L. japonica* possessed fucooidanase activity, of which twenty-eight strains, belonging mainly to *Alteromonas/Pseudoalteromonas*, could hydrolyze both fucooidans F<sub>1</sub> and F<sub>2</sub>, nine bacterial strains could hydrolyze fucooidan F<sub>2</sub>, and two strains, *Cytophaga* sp.

M3HL5/2 and *Alteromonas/Pseudoalteromonas* sp. M3HL14/2, could hydrolyze fucoidan F<sub>1</sub>. Strains M3HB2, M3HB9, M3HB14/1, M3HB15/3, and M3HL6/2 were the best producers of fucoidanases among the bacterial associates of the sea cucumber *A. japonicus*.

It should be noted that the fucoidanase activity of the most efficient fucoidan-degrading bacterial strains studied by us was only two–four times lower than that of *Vibrio* sp. 5, the proposed producer of fucoidanases [11]. However, the efficiency of this fucoidanase producer is much less than the efficiency of the known superproducers of laminarinases, amylases, cellulases, and other glycosidases.

The different activities of fucoidanases toward the three fucoidans studied are most likely due to the different structures of these substrates (see Table 1). It can be seen that fucoidans F<sub>1</sub>, F<sub>2</sub>, and F<sub>3</sub> considerably differ in molecular mass, carbohydrate composition, and the content of sulfate groups. Some of the fucoidanases studied were highly specific for particular fucoidans (Table 2). The relatively great number of bacterial strains that are able to hydrolyze fucoidan F<sub>3</sub> can be explained by the fact that this polysaccharide contains, in addition to fucose (42%), a large amount of galactose (40%). As a result, fucoidan F<sub>3</sub> can be hydrolyzed not only by fucoidanases but also by galactanases and galactosidases present in bacterial extracts.

The high percentage of fucoidan-degrading bacteria among the 78 bacterial strains studied (72% of strains isolated from brown seaweeds and 70.4% of strains isolated from sea cucumbers) indicates the nutritional importance of fucoidans for the bacterial associates of macrohydrobionts. The fucoidan-degrading ability may be part of the adaptive response of microorganisms colonizing seaweeds and various marine animals to the environment, through which these microorganisms satisfy their requirements for carbon, sulfur, and energy. The abundance of fucoidan-degrading marine bacteria and their low fucoidanase activity may be due to the specific role of fucoidans and fucoidanases in the metabolism of marine organisms.

## REFERENCES

- Kloareg, V. and Quatrano, R.S., Structure of the Cell Walls of Marine Algae and Ecophysiological Function of the Matrix Polysaccharides, *Oceanogr. Mar. Biol. Annu. Rev.*, 1988, vol. 26, pp. 259–315.
- Mulloy, V., Ribeiro, A., Alves, A., Vieira, R., and Mourao, P., Sulfated Fucans from Echinoderms Have a Regular Tetrasaccharide Repeating Unit Defined by Specific Patterns of Sulfation at the 0-2 and 0-4 Position, *J. Biol. Chem.*, 1994, vol. 269, no. 35, pp. 22113–22123.
- Glabe, C.G., Gabel, V., Vacquier, V.D., and Rosen, S.D., Carbohydrate Specificity of Sea Urchin Sperm Binding: a Cell Surface Lectin Mediating Sperm–Egg Adhesion, *J. Cell Biol.*, 1982, vol. 94, no. 1, pp. 123–128.
- Baba, M., Akajima, M., Schols, D., Pauwels, R., and Balzarini, J., Pentosan Polysulfate, a Sulfated Oligosaccharide, Is a Potent and Selective Anti-HIV Agent *In Vitro*, *Antiviral Res.*, 1988, vol. 9, no. 6, pp. 335–343.
- Nishino, T., Kiyohara, N., Yamaga, G., and Naguno, T., An Anticoagulant Fucoidan from the Brown Seaweed *Ecklonia kurome*, *Phytochemistry*, 1991, vol. 30, no. 2, pp. 535–539.
- Percival, E. and McDowell, R.H., *Chemistry and Enzymology of Marine Algal Polysaccharides*, New York: Academic, 1981, pp. 157–164.
- Usov, A.I., Smirnova, G.P., Bilan, M.I., and Shashkov, L.S., Algal Polysaccharides: The Brown Seaweed *Laminaria saccharine* as a Source of Fucoidan, *Bioorg. Khim.*, 1998, vol. 24, no. 6, pp. 437–445.
- Kitamura, K., Matsuo, M., and Yasui, T., Enzymic Degradation of Fucoidan by Fucoidanase from the Hepatopancreas of *Patinopecten yessoensis*, *Biosci. Biochem. Biotechnol.*, 1992, vol. 56, no. 3, pp. 490–494.
- Thanassi, N.M. and Nakada, G.I., Enzymic Degradation of Fucoidan by Enzymes from the Hepatopancreas of Abalone, *Arch. Biochem. Biophys.*, 1967, vol. 118, no. 1, pp. 172–177.
- Kitamura, K., Matsuo, M., and Yasui, T., Partial Purification and Characterization of an Enzyme Releasing 2-Sulfo- $\alpha$ -L-Fucopyranose From 2-Sulfo- $\alpha$ -L-Fucopyranosyl-(12)-Pyridylaminated Fucose from a Sea Urchin *Strongylocentrotus nudus*, *Biosci. Biochem. Biotechnol.*, 1996, vol. 60, no. 4, pp. 666–668.
- Furucawa, S., Fujikawa, T., Koga, D., and Ide, A., Purification and Some Properties of Exo-Type Fucoidanases from *Vibrio* sp. 5, *Biosci. Biochem. Biotechnol.*, 1992, vol. 56, no. 11, pp. 1829–1834.
- Ivanova, E.P., Bakunina, I.Yu., Nedashkovskaya, O.I., Gorshkova, N.M., Mikhailov, V.V., and Elyakova, L.A., Search for  $\alpha$ -Galactosidase Producers among Marine Bacteria of the Genus *Alteromonas*, *Prikl. Biokhim. Mikrobiol.*, 1996, vol. 32, no. 6, pp. 624–628.
- Zvyagintseva, T.N., Shirokova, N.I., and Elyakova, L.A., Structure of Laminarins from Some Brown Algae, *Bioorg. Khim.*, 1994, vol. 20, no. 12, pp. 1349–1358.
- Dubois, M., Gillers, K.A., Hamilton, J.K., Rebers, P.A., and Smith, F., Colorimetric Method for Determination of Sugars and Related Substances, *Anal. Chem.*, 1956, vol. 28, pp. 350–356.
- Nelson, T.E., A Photometric Adaptation of the Somogyi Method for the Determination of Glucose, *J. Biol. Chem.*, 1944, vol. 153, no. 1, pp. 375–381.
- Bakunina, I.Yu., Ivanova, E.P., Mikhailov, V.V., Nedashkovskaya, O.I., Gorshkova, N.M., and Parfenova, V.V., Distribution of  $\alpha$ -N-Galactosaminidases among Marine and Fresh-Water Microorganisms, *Mikrobiologiya*, 1994, vol. 63, no. 5, pp. 847–853.

17. Bakunina, I.Yu., Sova, V.V., Nedashkovskaya, O.I., Kul'man, R.A., Likhoshevstov, L.M., Martynova, M.A., Mikhailov, V.V., and Elyakova, L.A.,  $\alpha$ -Galactosidase from the Marine Bacterium *Pseudoalteromonas* sp. KMM 701, *Biokhimiya*, 1998, vol. 63, no. 10, pp. 1420–1427.
18. Warren, J., Microbial Hydrolysis of Polysaccharides, *Annu. Rev. Microbiol.*, 1996, vol. 50, pp. 183–212.
19. Laycock, R.A., The Detrital Food Chain Based on Seaweeds: 1. Bacteria Associated with the Surface of *Laminaria* Fronds, *Mar. Biol.*, 1974, vol. 25, no. 3, pp. 223–231.
20. Kong, M.K. and Chan, K., A Study of the Bacterial Flora Isolated from Marine Algae, *Bot. Mar.*, 1979, vol. 22, no. 2, pp. 83–97.
21. Jaffray, A.E., Anderson, R.J., and Coyne, V.E., Investigation of Bacterial Epiphytes of the Agar-Producing Red Seaweed *Gracilaria gracilis* (Stackhouse) Steentoft Irvine et Farnham from Saldanha Bay, South Africa and Luderitz, Namibia, *Bot. Mar.*, 1997, vol. 40, pp. 569–576.
22. Austin, B., Taxonomy of Marine Microorganisms, *Mar. Microbiol.*, Cambridge: Cambridge Univ. Press, 1988, p. 223.